

COCKLES

Co-Operation for Restoring Cockle Shellfisheries and its
Ecosystem-Services in the Atlantic Area

Modelling Larval Dispersal across the Atlantic Area


Sophie-Berenice Wilmes, Peter Robins, Ilaria Coscia, Sophie
Ward, Alice Goward-Brown, Luis Gimenez Noya, Enda O’Dea,
Elena Counago, Silvia Torres Lopez, and Shelagh Malham



FINAL VIRTUAL CONFERENCE

March 2021

Why are we interested in larval dispersal?




Connectivity is a measure of the rate of exchange of larvae between subpopulations

Important for:

- Determining which populations are more vulnerable
- Understanding oceanographic 'barriers'
- Management (MPA, commercial beds...)

Oceanography of the Atlantic Area

Monthly mean sea surface temperatures




IBI model PHYS_005_001 year 2018;
1/35° horiz. resolution

Ocean currents in the Atlantic Area

April - June

July - September



IBI model PHYS_005_001 year 2018; 15 m depth
1/35° horiz. resolution


Larval dispersal modelling

Focus on three areas:


- Full Atlantic Area
- Irish and Celtic Sea
- Galician Rias

Larval dispersal model:



- Virtual larvae released from cockle bed locations and advected for 35 days by ocean currents
- Last 10 days used for connectivity analysis
- Larval dispersal evaluated with releases from April to September
- Interannual variability considered where possible



Atlantic Area Connectivities



- Virtual larvae released from 51 sites
- April to September simulated
- 400 particles advected from each site per release day for 40 days
- Connectivity analysis based on 17,625,600 trajectories
- Last 10 days used for analysis




Atlantic Area connectivity


- Three distinct oceanographic units:
 1. Portugal, Spain, France up to Brittany
 2. English Channel
 3. Irish and Celtic Sea
- Divides more pronounced the deeper the larvae in the water column (more realistic?)
- Divides similar to patterns identified in the genetics analysis

Modelling larval dispersal in the Rias Baixas

Rias Baixas MOHID model



ROMS model NW Spain & Portugal




Merging of high and low resolution hydrodynamic models to achieve high resolution rias but also large enough domain

Particle tracking model setup:


- 200 virtual particles released each day
- April to September for the year 2018
- Advected for 35 days, with last 3 days used for connectivity analysis

Connectivity patterns Rias Baixas

April – September mean



01/06/2018 release



All Rias connected, but:

- Southern 3 sites show much stronger connectivities
- Fairly consistent pattern between months; weakest connectivities in August
- Weaker connectivity with Ria de Noia
- Implications for spread of Martilliosis?


Irish and Celtic Sea seascape genetics

Aims:

- Investigate population structure at finer scale resolution
- Understand drivers of genetic differentiation – environmental variables


Methods:

- Cockles sampled from 7 locations in 2010 and 2011
- Filtered dataset has 138 individuals and 1725 SNPs
- Larval dispersal modelling carried out using AMM NEMO model (1.5 km horizontal resolution)
- Link genetic and environmental data



Coscia et al., Fine-scale seascape genomics of an exploited marine species, the common cockle *Cerastoderma edule*, using a multimodelling approach, *Evolutionary Applications*, 2020.


Genetic substructure in the Irish and Celtic Seas



Distinct genetic substructure seen in neutral and outlier markers related to larval connectivity & environmental variables:

All sites connected, but distinct sub-units:

1. North Wales sites
2. Burry Inlet, Dyfi estuary & Gann
3. Irish sites and Gann



Mirrored in the genetics (neutral markers); outlier markers related to SSTs

Importance of larval dispersal for *C. edule* in the Atlantic Area

- Larval dispersal is an important part of the cockle's lifecycle during which subpopulations are able to connect with each other
- Understanding of larval dispersal important to determine vulnerable beds & thus inform management decisions
- Larval dispersal was investigated in 3 different areas:
 - Whole Atlantic Area: 3 distinct subgroups were demonstrated
 - Rias Baixas: All rias connected by larval dispersal but Ria de Noia less so
 - Irish Sea: 3 subgroups identified in the genetics related to larval dispersal
→ Work highlighted potential oceanographic barriers: headlands of Brittany and Cornwall, and Celtic Sea and Ushant Fronts
- Where is more work needed?
 - More knowledge on larval behaviour is required (ongoing experiments)
 - Links between estuaries and large-scale oceanography

Thanks for all of the fruitful discussions and enjoyable collaborations on the COCKLES project!